

## ZETA AT NEGATIVE ODD INTEGERS, A LA EULER

This writeup sketches (you may need to supply details) an argument due to Euler that partially establishes the functional equation of  $\zeta(s)$ . In particular, it leads to the formula

$$\zeta(1-k) = -\frac{B_k}{k} \quad \text{for even } k \geq 2.$$

Let  $t$  be a formal variable. Starting from the identity

$$t + t^2 + t^3 + t^4 + \dots = (t - t^2 + t^3 - t^4 + \dots) + 2(t^2 + t^4 + t^6 + t^8 + \dots),$$

apply the operator  $t \frac{d}{dt}$  (i.e., differentiation and then multiplication by  $t$ )  $n$  times to get

$$\begin{aligned} 1^n t + 2^n t^2 + 3^n t^3 + 4^n t^4 + \dots \\ = \left( t \frac{d}{dt} \right)^n \left( \frac{t}{1+t} \right) + 2^{n+1} (1^n t^2 + 2^n t^4 + 3^n t^6 + 4^n t^8 + \dots). \end{aligned}$$

Formally, when  $t = 1$  this is

$$\zeta(-n) = \left[ \left( t \frac{d}{dt} \right)^n \left( \frac{t}{1+t} \right) \right]_{t=1} + 2^{n+1} \zeta(-n),$$

giving a heuristic value for  $\zeta(-n)$ ,

$$\zeta(-n) = (1 - 2^{n+1})^{-1} \cdot \left[ \left( t \frac{d}{dt} \right)^n \left( \frac{t}{1+t} \right) \right]_{t=1}.$$

Thus for example, according to Euler,

$$1 + 1 + 1 + 1 + \dots = -\frac{1}{2} \quad \text{and} \quad 1 + 2 + 3 + 4 + \dots = -\frac{1}{12}.$$

Next, let  $t = e^X$  and note that  $t \frac{d}{dt} = \frac{d}{dX}$ . Now we have

$$(1 - 2^{n+1})\zeta(-n) = \left[ \frac{d^n}{dX^n} \left( \frac{e^X}{e^X + 1} \right) \right]_{X=0} \quad \text{for } n \in \mathbf{N},$$

giving the Taylor series

$$\frac{e^X}{e^X + 1} = \sum_{n=0}^{\infty} \frac{(1 - 2^{n+1})\zeta(-n)}{n!} X^n.$$

Thus the function of a complex variable

$$F(z) = \frac{e^{2\pi iz}}{e^{2\pi iz} + 1} = \sum_{n=0}^{\infty} \frac{(1 - 2^{n+1})\zeta(-n)(2\pi i)^n}{n!} z^n$$

generates (in the sense of generating function) the values  $\zeta(-n)$  for all natural numbers  $n$ .

Let  $G(z) = \pi \cot \pi z$ . Recall that like  $F$ ,  $G$  is both a fractional linear function of  $e^{2\pi iz}$  and a generating function for zeta,

$$G(z) = \pi i \frac{e^{2\pi iz} + 1}{e^{2\pi iz} - 1} = \frac{1}{z} - 2 \sum_{k=1}^{\infty} \zeta(2k) z^{2k-1}.$$

The fractional linear forms of  $F$  and  $G$  show that there must be a relation between them, and therefore between  $\zeta(1-k)$  and  $\zeta(k)$  for even  $k \geq 2$ . Indeed, the relation between  $F$  and  $G$  is

$$\frac{1}{\pi i} (G(z) - 2G(2z)) = -F(z) + F(-z),$$

and so equating coefficients gives

$$\zeta(1-k) = \frac{2\Gamma(k)}{(2\pi i)^k} \zeta(k) \quad \text{for even } k \geq 2.$$

The zeta-value that we have computed previously,

$$\zeta(k) = -\frac{1}{2} \cdot \frac{(2\pi i)^k B_k}{k!} \quad \text{for even } k \geq 2$$

now gives the identity promised at the beginning of the writeup,

$$\zeta(1-k) = -\frac{B_k}{k} \quad \text{for even } k \geq 2.$$

We want to symmetrize the previous two displays. A calculation shows that

$$\frac{\pi^{-\frac{1-k}{2}} \Gamma(\frac{1-k}{2})}{\pi^{-\frac{k}{2}} \Gamma(\frac{k}{2})} = \frac{1}{2} \cdot \frac{(2\pi i)^k}{(k-1)!}.$$

It follows that

$$\pi^{-k/2} \Gamma(\frac{k}{2}) \zeta(k) = \pi^{-(1-k)/2} \Gamma(\frac{1-k}{2}) \zeta(1-k) \quad \text{for even } k \geq 2.$$

This is a partial version of the *functional equation* of the zeta function. The full functional equation is

$$\pi^{-s/2} \Gamma(\frac{s}{2}) \zeta(s) = \pi^{-(1-s)/2} \Gamma(\frac{1-s}{2}) \zeta(1-s) \quad \text{for all } s \in \mathbf{C}.$$

This is usually written

$$\xi(s) = \xi(1-s) \quad \text{for all } s \in \mathbf{C}$$

where

$$\xi(s) = \pi^{-s/2} \Gamma(\frac{s}{2}) \zeta(s).$$

Euler's ideas here can be turned into a rigorous proof of the meromorphic continuation and the functional equation of  $\zeta$ , but the proof that we may give later in the course will not follow these lines. One comment to make for now is that the zeta function has an *Euler factorization*,

$$\zeta(s) = \prod_p (1 - p^{-s})^{-1} \quad \text{for } \operatorname{Re}(s) > 1,$$

where the product is taken over all primes  $p$ . From a modern perspective, each Euler factor  $(1 - p^{-s})^{-1}$  can be interpreted as an integral, taken in an environment particular to the prime  $p$ , and the extra factor  $\pi^{-s/2} \Gamma(s/2)$  in the symmetrized function  $\xi(s)$  is the same integral, but taken in an environment particular to a conceptually infinite prime.