

SIMPLICITY OF A_n

KEITH CONRAD

1. INTRODUCTION

A finite group is called *simple* when its only normal subgroups are the trivial subgroup and the whole group.

For instance, a finite group of prime size is simple, since it in fact has no non-trivial proper subgroups at all (normal or not). A finite abelian group G not of prime size, is not simple: let p be a prime factor of $\#G$, so G contains a subgroup of order p , which is a normal since G is abelian and is proper since $\#G > p$. Thus, the abelian finite simple groups are the groups of prime size.

When $n \geq 3$ the group S_n is not simple because it has a nontrivial normal subgroup A_n . But the groups A_n are simple, provided $n \geq 5$.

Theorem 1.1 (C. Jordan, 1875). *For $n \geq 5$, the group A_n is simple.*

The restriction $n \geq 5$ is optimal, since A_4 is not simple: it has a normal subgroup of size 4, namely $\{(1), (12)(34), (13)(24), (14)(23)\}$. The group A_3 is simple, since it has size 3, and the groups A_1 and A_2 are trivial.

We will give five proofs of Theorem 1.1. Section 2 includes some preparatory material and later sections give the proofs of Theorem 1.1. In the final section, we give a quick application of the simplicity of alternating groups and give references for further proofs not treated here.

2. PRELIMINARIES

We give two lemmas about alternating groups A_n for $n \geq 5$ and then two results on symmetric groups S_n for $n \geq 5$.

Lemma 2.1. *For $n \geq 3$, A_n is generated by 3-cycles. For $n \geq 5$, A_n is generated by permutations of type $(2, 2)$.*

Proof. That the 3-cycles generate A_n for $n \geq 3$ has been seen earlier in the course. To show permutations of type $(2, 2)$ generate A_n for $n \geq 5$, it suffices to write any 3-cycle (abc) in terms of such permutations. Pick $d, e \notin \{a, b, c\}$. Then note

$$(abc) = (ab)(de)(de)(bc).$$

□

The 3-cycles in S_n are all conjugate in S_n , since permutations of the same cycle type in S_n are conjugate. Are 3-cycles conjugate in A_n ? Not when $n = 4$: (123) and (132) are not conjugate in A_4 . But for $n \geq 5$ we do have conjugacy in A_n .

Lemma 2.2. *For $n \geq 5$, any two 3-cycles in A_n are conjugate in A_n .*

Proof. We show every 3-cycle in A_n is conjugate within A_n to (123). Let σ be a 3-cycle in A_n . It can be conjugated to (123) in S_n :

$$(123) = \pi\sigma\pi^{-1}$$

for some $\pi \in S_n$. If $\pi \in A_n$ we're done. Otherwise, let $\pi' = (45)\pi$, so $\pi' \in A_n$ and

$$\pi'\sigma\pi'^{-1} = (45)\pi\sigma\pi^{-1}(45) = (45)(123)(45) = (123).$$

□

Example 2.3. The 3-cycles (123) and (132) are not conjugate in A_4 . But in A_5 we have

$$(132) = \pi(123)\pi^{-1}$$

for $\pi = (45)(12) \in A_5$.

Most proofs of the simplicity of the groups A_n are based on Lemmas 2.1 and 2.2. The basic argument is this: show any non-trivial normal subgroup $N \triangleleft A_n$ contains a 3-cycle, so N contains every 3-cycle by Lemma 2.2, and therefore N is A_n by Lemma 2.1.

The next lemma will be used in our fifth proof of the simplicity of alternating groups.

Lemma 2.4. *For $n \geq 5$, the only nontrivial proper normal subgroup of S_n is A_n . In particular, the only subgroup of S_n with index 2 is A_n .*

Proof. The last statement follows from the first since any subgroup of index 2 is normal.

Let $N \triangleleft S_n$ with $N \neq \{(1)\}$. We will show $A_n \subset N$, so $N = A_n$ or S_n .

Pick $\sigma \in N$ with $\sigma \neq (1)$. That means there is an i with $\sigma(i) \neq i$. Pick $j \in \{1, 2, \dots, n\}$ so $j \neq i$ and $j \neq \sigma(i)$. Let $\tau = (ij)$. Then

$$\sigma\tau\sigma^{-1}\tau^{-1} = (\sigma(i) \sigma(j))(ij).$$

Since $\sigma(i) \neq i$ or j and $\sigma(i) \neq \sigma(j)$ (why?), the 2-cycles $(\sigma(i) \sigma(j))$ and (ij) are unequal, so their product is not the identity. That shows $\sigma\tau \neq \tau\sigma$.

Since $N \triangleleft S_n$, $\sigma\tau\sigma^{-1}\tau^{-1}$ lies in N . By construction, $\sigma(i) \neq i$ or j . If $\sigma(j) \neq i$ or j , then $(\sigma(i) \sigma(j))(ij)$ has type (2, 2). If $\sigma(j) = i$ or j , $(\sigma(i) \sigma(j))(ij)$ is a 3-cycle. Thus N contains a permutation of type (2, 2) or a 3-cycle. Since $N \triangleleft S_n$, N contains all permutations of type (2, 2) or all 3-cycles. In either case, this shows (by Lemma 2.1) that $N \supset A_n$. □

Remark 2.5. There is an analogue of Lemma 2.4 for the “countable” symmetric group S_∞ consisting of all permutations of $\{1, 2, 3, \dots\}$. A theorem of Schreier and Ulam (1933) says the only nontrivial proper normal subgroups of S_∞ are $\cup_{n \geq 1} S_n$ and $\cup_{n \geq 1} A_n$, which are the subgroup of permutations fixing all but a finite number of terms and its subgroup of even permutations.

Remark 2.6. From Lemma 2.4, any homomorphic image of S_n which is not an isomorphism has size 1 or 2. In particular, there is no surjective homomorphism $S_n \rightarrow \mathbf{Z}/(m)$ for $m > 2$.

Theorem 2.7. *For $n \geq 5$, any proper subgroup of S_n other than A_n has index at least n . Moreover, any subgroup of index n is isomorphic to S_{n-1} .*

Proof. Let H be a proper subgroup of S_n other than A_n , and let $m > 1$ be the index of H in S_n . We want to show $m \geq n$. Assume $m < n$. The left multiplication action of S_n on S_n/H gives a group homomorphism

$$\varphi: S_n \rightarrow \text{Sym}(S_n/H) \cong S_m.$$

By hypothesis, $m < n$, so φ is not injective. Let K be the kernel of φ , so $K \subset H$ and K is non-trivial. Since $K \triangleleft S_n$, Lemma 2.4 says $K = A_n$ or S_n . Since $K \subset H$, we get $H = A_n$ or S_n , which contradicts our initial assumption about H . Therefore $m \geq n$.

Now let H be a subgroup of S_n with index n . Consider the left multiplication action of S_n on S_n/H . This is a homomorphism $\ell: S_n \rightarrow \text{Sym}(S_n/H)$. Since S_n/H has size n , $\text{Sym}(S_n/H)$ is isomorphic to S_n . The kernel of ℓ is a normal subgroup of S_n which lies in H (why?). Therefore the kernel has index at least n in S_n . Since the only normal subgroups of S_n are 1, A_n , and S_n , the kernel of ℓ is trivial, so ℓ is an isomorphism. What is the image $\ell(H)$ in $\text{Sym}(S_n/H)$? Since $gH = H$ if and only if $g \in H$, $\ell(H)$ is the group of permutations of S_n/H which fixes the ‘‘point’’ H in S_n/H . The subgroup fixing a point in a symmetric group isomorphic to S_n is isomorphic to S_{n-1} . Therefore $H \cong \ell(H) \cong S_{n-1}$. \square

Theorem 2.7 is false for $n = 4$: S_4 contains the dihedral group of size 8 as a subgroup of index 3. An analogue of Theorem 2.7 for alternating groups will be given in Section 8; its proof uses the simplicity of alternating groups.

Corollary 2.8. *Let F be a field. If $f \in F[X_1, \dots, X_n]$ and $n \geq 5$, the number of different polynomials we get from f by permuting its variables is either 1, 2, or at least n .*

Proof. Letting S_n act on $F[X_1, \dots, X_n]$ by permutations of the variables, the polynomials we get by permuting the variables of f is the S_n -orbit of f . The size of this orbit is $[S_n : H]$, where $H = \text{Stab}_f = \{\sigma \in S_n : \sigma f = f\}$. By Theorem 2.7, this index is either 1, 2, or at least n . \square

3. FIRST PROOF

Our first proof of Theorem 1.1 is based on the one in [2, pp. 149–150].

We begin by showing A_5 is simple.

Theorem 3.1. *The group A_5 is simple.*

Proof. We want to show the only normal subgroups of A_5 are $\{(1)\}$ and A_5 . This will be done in two ways.

Our first method involves counting the sizes of the conjugacy classes. There are 5 conjugacy classes in A_5 , with representatives and sizes as indicated in the following table.

Rep.	(1)	(12345)	(21345)	(12)(34)	(123)
Size	1	12	12	15	20

If A_5 has a normal subgroup N , then N is a union of conjugacy classes – including $\{(1)\}$ – whose total size divides 60. However, no sum of the above numbers which includes 1 is a factor of 60 except for 1 and 60. Therefore N is trivial or A_5 .

For the second proof, let $N \triangleleft A_5$ with $\#N > 1$. We will show N contains a 3-cycle. It follows that $N = A_n$ by Lemmas 2.1 and 2.2.

Pick $\sigma \in N$ with $\sigma \neq (1)$. The cycle structure of σ is (abc) , $(ab)(cd)$, or $(abcde)$, where different letters represent different numbers. Since we want to show N contains a 3-cycle, we may suppose σ has the second or third cycle type. In the second case, N contains

$$((abe)(ab)(cd)(abe)^{-1})(ab)(cd) = (be)(cd)(ab)(cd) = (aeb).$$

In the third case, N contains

$$((abc)(abcde)(abc)^{-1})(abcde)^{-1} = (adebc)(aedcb) = (abd).$$

Therefore N contains a 3-cycle, so $N = A_5$. \square

Lemma 3.2. *When $n \geq 5$, any $\sigma \neq (1)$ in A_n has a conjugate $\sigma' \neq \sigma$ such that $\sigma(i) = \sigma'(i)$ for some i .*

For example, if $\sigma = (12345)$ in A_5 then $\sigma' = (345)\sigma(345)^{-1} = (12453)$ has the same value at $i = 1$ as σ does.

Proof. Let σ be a non-identity element of A_n . Let r be the longest length of a disjoint cycle in σ . Relabelling, we may write

$$\sigma = (12 \dots r)\pi,$$

where $(12 \dots r)$ and π are disjoint.

If $r \geq 3$, let $\tau = (345)$ and $\sigma' = \tau\sigma\tau^{-1}$. Then $\sigma(1) = 2, \sigma'(1) = 2, \sigma(2) = 3$, and $\sigma'(2) = 4$. Thus $\sigma' \neq \sigma$ and both take the same value at 1.

If $r = 2$, then σ is a product of disjoint transpositions. If there are at least 3 disjoint transpositions involved, then $n \geq 6$ and we can write $\sigma = (12)(34)(56)(\dots)$ after relabelling. Let $\tau = (12)(35)$ and $\sigma' = \tau\sigma\tau^{-1}$. Then $\sigma(1) = 2, \sigma'(1) = 2, \sigma(3) = 4$, and $\sigma'(3) = 6$. Again, we see $\sigma' \neq \sigma$ and σ and σ' have the same value at 1.

If $r = 2$ and σ is a product of 2 disjoint transpositions, write $\sigma = (12)(34)$ after relabelling. Let $\tau = (132)$ and $\sigma' = \tau\sigma\tau^{-1} = (13)(24)$. Then $\sigma' \neq \sigma$ and they both fix 5. \square

Now we prove Theorem 1.1.

Proof. We may suppose $n \geq 6$, by Theorem 3.1. For $1 \leq i \leq n$, let A_n act in the natural way on $\{1, 2, \dots, n\}$ and let $H_i \subset A_n$ be the subgroup fixing i , so $H_i \cong A_{n-1}$. By induction, each H_i is simple. Note each H_i contains a 3-cycle (build out of 3 numbers other than i).

Let $N \triangleleft A_n$ be a nontrivial normal subgroup. We want to show $N = A_n$. Pick $\sigma \in N$ with $\sigma \neq \{(1)\}$. By Lemma 3.2, there is a conjugate σ' of σ such that $\sigma' \neq \sigma$ and $\sigma(i) = \sigma'(i)$ for some i . Since N is normal in A_n , $\sigma' \in N$. Then $\sigma^{-1}\sigma'$ is a non-identity element of N which fixes i , so $N \cap H_i$ is a non-trivial subgroup of H_i . It is also a normal subgroup of H_i since $N \triangleleft A_n$. Since H_i is simple, $N \cap H_i = H_i$. Therefore $H_i \subset N$. Since H_i contains a 3-cycle, N contains a 3-cycle and we are done.

Alternatively, we can show $N = A_n$ when $N \cap H_i$ is non-trivial for some i as follows. As before, since $N \cap H_i$ is a non-trivial normal subgroup of H_i , $H_i \subset N$. Without referring to 3-cycles, we instead note that the different H_i 's are conjugate subgroups of A_n : $\sigma H_i \sigma^{-1} = H_{\sigma(i)}$ for $\sigma \in A_n$. Since $N \triangleleft A_n$ and N contains H_i , N contains every $H_{\sigma(i)}$ for all $\sigma \in A_n$. Since $\sigma(i)$ can be any element of A_n as σ varies in A_n , N contains every H_i . Any permutation of type $(2, 2)$ is in some H_i since $n \geq 5$, so N contains all permutations of type $(2, 2)$. Every permutation in A_n is a product of permutations of type $(2, 2)$, so $N \supset A_n$. Therefore $N = A_n$. \square

4. SECOND PROOF

Our next proof is taken from [6, p. 108]. It does not use induction on n , but we do need to know A_6 is simple at the start.

Theorem 4.1. *The group A_6 is simple.*

Proof. We follow the first method of proof of Theorem 3.1. Here is the table of conjugacy classes in A_6 .

Rep.	(1)	(123)	(123)(456)	(12)(34)	(12345)	(23456)	(1234)(56)
Size	1	40	40	45	72	72	90

A tedious check shows no sum of these sizes, which includes 1, is a factor of $6!/2$ except for the sum of all the terms. Therefore the only non-trivial normal subgroup of A_6 is A_6 . \square

Now we prove the simplicity of A_n for larger n by reducing directly to the case of A_6 .

Proof. Since A_5 and A_6 are known to be simple by Theorems 3.1 and 4.1, pick $n \geq 7$ and let $N \triangleleft A_n$ be a non-trivial subgroup. We will show N contains a 3-cycle.

Let σ be a non-identity element of N . It moves some number. By relabelling, we may suppose $\sigma(1) \neq 1$. Let $\tau = (ijk)$, where i, j, k are not 1 and $\sigma(1) \in \{i, j, k\}$. Then $\tau\sigma\tau^{-1}(1) = \tau(\sigma(1)) \neq \sigma(1)$, so $\tau\sigma\tau^{-1} \neq \sigma$. Let $\varphi = \tau\sigma\tau^{-1}\sigma^{-1}$, so $\varphi \neq (1)$. Writing

$$\varphi = (\tau\sigma\tau^{-1})\sigma^{-1},$$

we see $\varphi \in N$. Now write

$$\varphi = \tau(\sigma\tau^{-1}\sigma^{-1}),$$

Since τ^{-1} is a 3-cycle, $\sigma\tau^{-1}\sigma^{-1}$ is also a 3-cycle. Therefore φ is a product of two 3-cycles, so φ moves at most 6 numbers in $\{1, 2, \dots, n\}$. Let H be the copy of A_6 inside A_n corresponding to the even permutations of those 6 numbers (possibly augmented to 6 arbitrarily if in fact φ moves fewer numbers). Then $N \cap H$ is non-trivial (it contains φ) and it is a normal subgroup of H . Since $H \cong A_6$, which is simple, $N \cap H = H$. Thus $H \subset N$, so N contains a 3-cycle. \square

5. THIRD PROOF

Our next proof is by induction, and uses conjugacy classes instead of Lemma 3.2. It is based on [9, p. 5].

Lemma 5.1. *If $n \geq 6$ then every non-trivial conjugacy class in S_n and A_n has at least n elements.*

The lower bound n in Lemma 5.1 is actually quite weak as n grows. But it shows that the size of each non-trivial conjugacy class in S_n and A_n grows with n .

Proof. For $n \geq 6$, pick $\sigma \in S_n$ with $\sigma \neq (1)$. We want to look at the conjugacy class of σ in S_n , and if $\sigma \in A_n$ we also want to look at the conjugacy class of σ in A_n , and our goal in both cases is to find at least n elements in the conjugacy class.

Case 1: The disjoint cycle decomposition of σ includes a cycle with length greater than 2. Without loss of generality, $\sigma = (123\dots)\dots$

For $3 \leq k \leq n$, fix a choice of $\ell \notin \{1, 2, 3, k\}$ (which is possible since $n \geq 5$) and let $\alpha_k = (2k\ell)$ and $\beta_k = (3k\ell)$. Then $\alpha_k\sigma\alpha_k^{-1}$ has the effect $1 \rightarrow 1 \rightarrow 2 \rightarrow k$ and $\beta_k\sigma\beta_k^{-1}$ has the effect $1 \rightarrow 1 \rightarrow 2 \rightarrow 2$ and $2 \rightarrow 2 \rightarrow 3 \rightarrow k$. This tells us that the conjugates

$$\alpha_3\sigma\alpha_3^{-1}, \dots, \alpha_n\sigma\alpha_n^{-1}, \beta_3\sigma\beta_3^{-1}, \dots, \beta_n\sigma\beta_n^{-1}$$

are all different from each other: the conjugates by the α 's have different effects on 1, the conjugates by the β 's have different effects on 2, and a conjugate by an α is not a conjugate by a β since they have different effects on 1. Since these conjugates are different, the number of conjugates of σ is at least $2(n-2) > n$. Because α_k and β_k are 3-cycles, if $\sigma \in A_n$ then these conjugates are in the A_n -conjugacy class of σ .

Case 2: The disjoint cycle decomposition of σ only has cycles with length 1 or 2. Therefore without loss of generality σ is a transposition or a product of at least 2 disjoint transpositions.

If σ is a transposition, then its S_n -conjugacy class is the set of all transpositions (ij) where $1 \leq i < j \leq n$, and the number of these permutations is $\binom{n}{2} = \frac{n^2-n}{2}$, which is greater than n for $n \geq 6$.

If σ is a product of at least 2 disjoint transpositions, then without loss of generality $\sigma = (12)(34)\dots$, where the terms in \dots don't involve 1, 2, 3, or 4.

For $5 \leq k \leq n$, let $\alpha_k = (12)(3k)$, $\beta_k = (13)(2k)$, and $\gamma_k = (1k)(23)$. Then $\alpha_k \sigma \alpha_k^{-1}$ has the effect

$$1 \rightarrow 2 \rightarrow 1 \rightarrow 2, \quad 2 \rightarrow 1 \rightarrow 2 \rightarrow 1, \quad k \rightarrow 3 \rightarrow 4 \rightarrow 4,$$

$\beta_k \sigma \beta_k^{-1}$ has the effect

$$1 \rightarrow 3 \rightarrow 4 \rightarrow 4, \quad 3 \rightarrow 1 \rightarrow 2 \rightarrow k, \quad k \rightarrow 2 \rightarrow 1 \rightarrow 3,$$

and $\gamma_k \sigma \gamma_k^{-1}$ has the effect

$$2 \rightarrow 3 \rightarrow 4 \rightarrow 4, \quad 3 \rightarrow 2 \rightarrow 1 \rightarrow k, \quad k \rightarrow 1 \rightarrow 2 \rightarrow 3.$$

The conjugates of σ by the α 's are different from each other since they take different elements to 4, the conjugates of σ by the β 's are different from each other since they take different elements to 3, and the conjugates of σ by the γ 's are different from each other since they take different elements to 3. Conjugates of σ by an α and a β are different since they send 1 to different places, conjugates of σ by an α and a γ are different since they send 2 to different places, and conjugates of σ by a β and a γ are different since they send different elements to 4 (1 for the β 's and 2 for the γ 's). In total the number of conjugates of σ we have written down (which are all conjugates by 3-cycles, hence they are conjugates in A_n if $\sigma \in A_n$) is $3(n-4)$, and $3(n-4) \geq n$ if $n \geq 6$. \square

Now we prove Theorem 1.1.

Proof. We argue by induction on n , the case $n = 5$ having already been settled by Theorem 3.1. Say $n \geq 6$. Let $N \triangleleft A_n$ with $N \neq \{(1)\}$. Since N is normal and non-trivial, it contains non-identity conjugacy classes in A_n . By Lemma 5.1, any non-identity conjugacy class in A_n has size at least n when $n \geq 6$. Therefore, by counting the trivial conjugacy class and a non-trivial conjugacy class in N , we see $\#N \geq n + 1$.

Using a wholly different argument, we now show that $\#N \leq n$ if $N \neq A_n$, which will be a contradiction. Pick $1 \leq i \leq n$. Let $H_i \subset A_n$ be the subgroup fixing i , so $H_i \cong A_{n-1}$. In particular, H_i is a simple group by induction. Notice each H_i contains a 3-cycle.

The intersection $N \cap H_i$ is a normal subgroup of H_i , so simplicity of H_i implies $N \cap H_i$ is either $\{(1)\}$ or H_i . If $N \cap H_i = H_i$ for some i , then $H_i \subset N$. Since H_i contains a 3-cycle, N does as well, so $N = A_n$ by Lemmas 2.1 and 2.2. (This part resembles part of our first proof of simplicity of A_n , but we will now use Lemma 5.1 instead of Lemma 3.2 to show the possibility that $N \cap H_i = \{(1)\}$ for all i is absurd.)

Suppose $N \neq A_n$. Then, by the previous paragraph, $N \cap H_i = \{(1)\}$ for all i . Therefore each $\sigma \neq (1)$ in N acts on $\{1, 2, \dots, n\}$ without fixed points (otherwise σ would be a non-identity element in some $N \cap H_i$). That implies each $\sigma \neq (1)$ in N is completely determined by the value $\sigma(1)$: if $\tau \neq (1)$ is in N and $\sigma(1) = \tau(1)$, then $\sigma\tau^{-1} \in N$ fixes 1, so $\sigma\tau^{-1}$ is the identity, so $\sigma = \tau$.

There are only $n - 1$ possible values for $\sigma(1) \in \{2, 3, \dots, n\}$, so $N - \{(1)\}$ has size at most $n - 1$, hence $\#N \leq n$. We already saw from Lemma 5.1 that $\#N \geq n + 1$, so we have a contradiction. \square

6. FOURTH PROOF

Our next proof, based on [3, p. 50], is very computational.

Proof. Let $N \triangleleft A_n$ be a non-trivial normal subgroup. We will show N contains a 3-cycle.

Pick $\sigma \in N$, $\sigma \neq (1)$. Write

$$\sigma = \pi_1 \pi_2 \cdots \pi_k,$$

where the π_j 's are disjoint cycles. In particular, they *commute*, so we can relabel them at our convenience. Eliminate any 1-cycles from the product.

Case 1: Some π_i has length at least 4. Relabelling, we can write

$$\pi_1 = (12 \cdots r)$$

with $r \geq 4$. Let $\varphi = (123)$. Then $\varphi \sigma \varphi^{-1} \in N$ and

$$\begin{aligned} \varphi \sigma \varphi^{-1} &= \varphi \pi_1 \varphi^{-1} \pi_2 \cdots \pi_k \\ &= \varphi \pi_1 \varphi^{-1} \pi_1^{-1} \sigma \\ &= (123)(123 \cdots r)(132)(r \cdots 21)\sigma \\ &= (124)\sigma, \end{aligned}$$

so $(124) = \varphi \sigma \varphi^{-1} \sigma^{-1} \in N$.

Case 2: Each π_i has length ≤ 3 , and at least two have length 3 (so $n \geq 6$). Without loss of generality, $\pi_1 = (123)$ and $\pi_2 = (456)$. Let $\varphi = (124)$. Then

$$\begin{aligned} \varphi \sigma \varphi^{-1} &= \varphi \pi_1 \pi_2 \varphi^{-1} \pi_3 \cdots \pi_k \\ &= \varphi \pi_1 \pi_2 \varphi^{-1} \pi_2^{-1} \pi_1^{-1} \sigma \\ &= (124)(123)(456)(142)(465)(132)\sigma \\ &= (12534)\sigma, \end{aligned}$$

so $\varphi \sigma \varphi^{-1} \sigma^{-1} = (12534) \in N$. Now run through Case 1 with this 5-cycle to find a 3-cycle in N .

Case 3: Exactly one π_i has length 3, and the rest have length ≤ 2 . Without loss of generality, $\pi_1 = (123)$ and the other π_i 's are 2-cycles. Then $\sigma^2 = \pi_1^2$ is in N , and $\pi_1^2 = (132)$.

Case 4: All π_i 's are 2-cycles, so necessarily $k > 1$. Write $\pi_1 = (12)$ and $\pi_2 = (34)$. Let $\varphi = (123)$. Then

$$\begin{aligned} \varphi \sigma \varphi^{-1} &= \varphi \pi_1 \pi_2 \varphi^{-1} \pi_3 \cdots \pi_k \\ &= \varphi \pi_1 \pi_2 \varphi^{-1} \pi_2^{-1} \pi_1^{-1} \sigma \\ &= (123)(12)(34)(132)(34)(12)\sigma \\ &= (13)(24)\sigma, \end{aligned}$$

so

$$\varphi \sigma \varphi^{-1} \sigma^{-1} = (13)(24) \in N.$$

Let $\psi = (135)$. Then

$$\begin{aligned} (13)(24)\psi(13)(24)\psi^{-1} &= (13)(24)(135)(13)(24)(153) \\ &= (13)(135)(13)(153) \\ &= (135), \end{aligned}$$

so N contains a 3-cycle. □

7. FIFTH PROOF

Our final proof is taken from [8, p. 295].

Let $N \triangleleft A_n$ with N not $\{(1)\}$ or A_n . We will study N as a subgroup of S_n . By Lemma 2.4, N is not a normal subgroup of S_n . This means the normalizer of N inside S_n is a proper subgroup, which contains A_n , so

$$(7.1) \quad A_n = N_{S_n}(N).$$

For any transposition τ in S_n , $\tau \notin N_{S_n}(N)$ by (7.1), so $\tau N \tau^{-1} \neq N$. Since $N \triangleleft A_n$ and $\tau N \tau^{-1}$ is a subgroup of A_n , the product set $N \cdot \tau N \tau^{-1}$ is a subgroup of A_n . We have the chain of inclusions

$$N \cap \tau N \tau^{-1} \subset N \subset N \cdot \tau N \tau^{-1} \subset A_n,$$

where the first and second are strict.

We will now show, for any transposition τ in S_n , that

$$(7.2) \quad N \cap \tau N \tau^{-1} \triangleleft S_n, \quad N \cdot \tau N \tau^{-1} \triangleleft S_n.$$

The proof of (7.2) is a bit tedious, so first let's see why (7.2) leads to a contradiction.

It follows from (7.2) and Lemma 2.4 that

$$(7.3) \quad N \cap \tau N \tau^{-1} = \{(1)\}, \quad N \cdot \tau N \tau^{-1} = A_n$$

for any transposition τ in S_n . By (7.3), $\#A_n = \#N \cdot \#(\tau N \tau^{-1}) = (\#N)^2$, so $n! = 2(\#N)^2$. This tells us $\#N$ must be even, so N has an element, say σ , of order 2. Then σ is a product of disjoint 2-cycles. There is a transposition ρ in S_n which commutes with σ : just take for ρ one of the transpositions in the disjoint cycle decomposition of σ . Then

$$\sigma = \rho \sigma \rho^{-1} \in N \cap \rho N \rho^{-1}.$$

From (7.3), using ρ for the arbitrary τ there, $N \cap \rho N \rho^{-1}$ is trivial, so we have a contradiction. (As another way of reaching a contradiction from the equation $n! = 2(\#N)^2$, we can use Bertrand's postulate – proved by Chebyshev – that there is always a prime strictly between m and $2m$ for any $m > 1$. That means, taking $m = n!/4$, the ratio $n!/2$ can't be a perfect square.)

It remains to check the two conditions in (7.2). In both cases, we show the subgroups are normalized by A_n and by τ , so the normalizer contains $\langle A_n, \tau \rangle = S_n$.

First consider $N \cap \tau N \tau^{-1}$. It is clearly normalized by τ . Now pick any $\pi \in A_n$. Then $\pi N \pi^{-1} = N$ since $N \triangleleft A_n$, and

$$(7.4) \quad \pi(\tau N \tau^{-1})\pi^{-1} = \tau(\tau^{-1}\pi\tau)N(\tau^{-1}\pi^{-1}\tau)\tau^{-1} = \tau N \tau^{-1}$$

since $\tau^{-1}\pi\tau \in A_n$. Therefore

$$\pi(N \cap \tau N \tau^{-1})\pi^{-1} = \pi N \pi^{-1} \cap \pi \tau N \tau^{-1} \pi^{-1} = N \cap \tau N \tau^{-1},$$

so A_n normalizes $N \cap \tau N \tau^{-1}$.

Now we look at $N \cdot \tau N \tau^{-1}$. Pick an element of this product, say

$$\sigma = \sigma_1 \tau \sigma_2 \tau^{-1},$$

where $\sigma_1, \sigma_2 \in N$. Then, since $N \triangleleft A_n$,

$$\tau \sigma \tau^{-1} = \tau \sigma_1 \tau \sigma_2 \tau^{-2} = \tau \sigma_1 \tau \sigma_2 \in \tau N \tau^{-1} \cdot N = N \cdot \tau N \tau^{-1},$$

which shows τ normalizes $N \cdot \tau N \tau^{-1}$.

Now pick any $\pi \in A_n$. To see π normalizes $N \cdot \tau N \tau^{-1}$, pick σ as before. Then

$$\pi \sigma \pi^{-1} = \pi \sigma_1 \pi^{-1} \cdot \pi (\tau \sigma_2 \tau^{-1}) \pi^{-1}.$$

The first factor $\pi \sigma_1 \pi^{-1}$ is in N since $N \triangleleft A_n$. The second factor is in $\pi \tau N \tau^{-1} \pi^{-1}$, which equals $\tau N \tau^{-1}$ by (7.4).

8. CONCLUDING REMARKS

The standard counterexample to the converse of Lagrange's theorem is A_4 : it has size 12 but no subgroup of index 2. For $n \geq 5$, the groups A_n also have no subgroup of index 2, since any index-2 subgroup of a group is normal and A_n is simple.

In fact, something stronger is true.

Corollary 8.1. *For $n \geq 5$, any proper subgroup of A_n has index at least n .*

This is an analogue of Theorem 2.7.

Proof. Let H be a proper subgroup of A_n , with index $m > 1$. Consider the left multiplication action of A_n on A_n/H . This gives a group homomorphism

$$\varphi: A_n \rightarrow \text{Sym}(A_n/H) \cong S_m.$$

Let K be the kernel of φ , so $K \subset H$ (why?) and $K \triangleleft A_n$. By simplicity of A_n , K is trivial. Therefore A_n injects into S_m , so $(n!/2) | m!$, which implies $n \leq m$. \square

The lower bound of n is sharp since $[A_n : A_{n-1}] = n$. Corollary 8.1 is false for $n = 4$: A_4 has a subgroup of index 3.

Remark 8.2. What the proof of Corollary 8.1 shows more generally is that if G is a finite simple group and H is a subgroup with index $m > 1$, then there is an embedding of G into S_m , so $\#G | m!$. With G fixed, this divisibility relation puts a lower bound on the index of any proper subgroup of G .

A reader who wants to see more proofs that A_n is simple for $n \geq 5$ can look at [4, pp. 247–248] or [5, pp. 32–33] for another way of showing a non-trivial normal subgroup contains a 3-cycle, or at [1, §1.7] or [7, pp. 295–296] for a proof based on the theory of highly transitive permutation groups.

REFERENCES

- [1] N. L. Biggs, A. T. White, "Permutation Groups and Combinatorial Structures," Cambridge Univ. Press, Cambridge, 1979.
- [2] D. Dummit, R. Foote, "Abstract Algebra," 3rd ed., J. Wiley, 2004.
- [3] T. Hungerford, "Algebra," Springer-Verlag, New York, 1980.
- [4] N. Jacobson, "Basic Algebra I," 2nd ed., W. H. Freeman, New York, 1985.
- [5] S. Lang, "Algebra," revised 3rd ed., Springer-Verlag, New York, 2002.
- [6] J. Rotman, "Advanced Modern Algebra," Prentice-Hall, Upper Saddle River, 2002.
- [7] W. R. Scott, "Group Theory," Dover, New York, 1987.
- [8] M. Suzuki, "Group Theory I," Springer-Verlag, Berlin, 1982.
- [9] R. Wilson, "Alternating Groups," at <http://www.maths.qmw.ac.uk/~raw/fsgs.html>.