

**MATH 341: TOPICS IN GEOMETRY
HOMEWORK DUE FRIDAY WEEK 4**

Problem 1. This is a continuation of Problem 5 from HW2, and you may freely use the conclusions of that problem here.

Let $\triangle ABC$ be a triangle in the Euclidean plane. Let A^* denote the point in the plane such that $\triangle A^*BC$ is equilateral and such that A and A^* lie on opposite sides of the line through B and C . Let θ_A denote the rotation about the circumcenter¹ O_A of $\triangle A^*BC$ which takes B to C . Similarly define θ_C and θ_B .

- (a) Show that $\theta_C\theta_B\theta_A = \text{id}$. *Hint:* Show that $\theta_C\theta_B\theta_A$ is a rotation by angle 0.
- (b) Extend the notation above in the natural way and then show that the circumcenters $O_A, O_B,$ and O_C form an equilateral triangle. *Hint:* Calculate $\theta_B\theta_A$.

Problem 2. Let E denote an n -dimensional Euclidean space and let k and ℓ be lines through the origin in E . Define the *acute angle* $\angle(k, \ell) \in [0, \pi/2]$ between k and ℓ by the formula

$$\angle(k, \ell) = \arccos \left| \langle \hat{k}, \hat{\ell} \rangle \right|$$

where \hat{k} and $\hat{\ell}$ are unit vectors generating k and ℓ , respectively.

- (a) Show that the acute angle defines a metric on *projective space* $P(E)$, the set of lines through the origin in E . *Hint:* Interpret $P(E)$ as the orbit space for the action of the antipodal map on the sphere $S(E)$, and use what we have developed about spherical geometry.
- (b) Show that there is a unique geodesic passing through any two distinct points of $P(E)$.

Problem 3. The *torus* is the set $S^1 \times S^1$ endowed with the product metric from Problem 4 of the previous problem set (where $S^1 = S(\mathbb{R}^2)$ is endowed with the standard spherical metric). The canonical map $\chi: \mathbb{R} \rightarrow S^1, \theta \mapsto (\cos \theta, \sin \theta)$ induces a map

$$\begin{aligned} \nu: \mathbb{R}^2 &\longrightarrow S^1 \times S^1 \\ (\theta, \varphi) &\longmapsto (\chi(\theta), \chi(\varphi)). \end{aligned}$$

- (a) For points $p, q \in S^1 \times S^1$, show that

$$d(p, q) = \inf_{\{(P, Q) \in \mathbb{R}^2 \times \mathbb{R}^2 \mid p = \nu(P), q = \nu(Q)\}} d(P, Q).$$

- (b) Show that ν is a *local isometry*: there exists a constant $r > 0$ such that for all $P \in \mathbb{R}^2$ the open disk with center P and radius r is mapped bijectively by ν onto the “metric disk” $\{q \in S^1 \times S^1 \mid d(q, \nu(P)) < r\}$ in $S^1 \times S^1$.
- (c) Show that an affine line in \mathbb{R}^2 is mapped by ν onto a geodesic in $S^1 \times S^1$ and that all geodesics in $S^1 \times S^1$ have this form.

¹The *circumcenter* of a triangle is the center of the (unique) circle passing through the vertices of the triangle.