

Math 322

April 11, 2022

Presentations

Presentation proposals due next Monday.

Index of a vector field on a manifold

- ▶ oriented 2-dimensional manifold (surface)
- ▶ vector field on S
- ▶ index of a critical point on a manifold

Poincaré-Hopf index theorem

Theorem. Let f be a vector field on a compact surface S of genus g , Suppose the critical points of f are p_1, \dots, p_k . Then

$$\sum_{i=1}^k I_f(S) = 2 - 2g.$$

Poincaré-Hopf index theorem

Theorem. Let f be a vector field on a compact surface S of genus g , Suppose the critical points of f are p_1, \dots, p_k . Then

$$\sum_{i=1}^k I_f(S) = 2 - 2g.$$


Proof. Case 1: $S = S^2$.

Poincaré-Hopf index theorem

Theorem. Let f be a vector field on a compact surface S of genus g . Suppose the critical points of f are p_1, \dots, p_k . Then

$$\sum_{i=1}^k I_f(S) = 2 - 2g.$$

Proof. Case 1: $S = S^2$.




Poincaré-Hopf index theorem

Theorem. Let f be a vector field on a compact surface S of genus g . Suppose the critical points of f are p_1, \dots, p_k . Then

$$\sum_{i=1}^k I_f(S) = 2 - 2g.$$

Proof. Case 1: $S = S^2$.




Poincaré-Hopf index theorem

Theorem. Let f be a vector field on a compact surface S of genus g . Suppose the critical points of f are p_1, \dots, p_k . Then

$$\sum_{i=1}^k I_f(S) = 2 - 2g.$$

Proof. Case 1: $S = S^2$.




Poincaré-Hopf index theorem

Theorem. Let f be a vector field on a compact surface S of genus g . Suppose the critical points of f are p_1, \dots, p_k . Then


$$\sum_{i=1}^k I_f(S) = 2 - 2g.$$

Proof. Case 1: $S = S^2$.



Poincaré-Hopf index theorem

General case:



Poincaré-Hopf index theorem

General case:

