

MATHEMATICS 311: COMPLEX ANALYSIS — ASSIGNMENT 6

**Reading:** Marsden, section 3.3.

**Problems:** 1. Let  $f(z)$  be analytic in the entire complex plane  $\mathbb{C}$ , and set  $M(r) = \sup_{|z|=r} |f(z)|$ . Show that  $M(r_1) \leq M(r_2)$  whenever  $r_1 < r_2$ , and determine when equality can occur.

2. Find all expansions in powers of  $z$  of the function

$$f(z) = \frac{1}{z^2(z^2 + 1)(z^2 + 9)}.$$

(Note that  $z^2$  is already a power of  $z$ . Use partial fractions and then the handy formulas from class.)

3. Find all expansions in powers of  $z - 1$  and in powers of  $z + 1/2$  of the function

$$f(z) = \frac{1}{(2z + 1)(z - 1)}.$$

(There is no need to use partial fractions on this problem.)

4. Suppose  $f(z)$  is analytic in the strip  $a < y < b$  (where  $z = x + iy$ ) and satisfies  $f(z + 1) = f(z)$ . Show that  $f(z)$  has a complex Fourier expansion

$$f(z) = \sum_{n=-\infty}^{\infty} a_n e^{2\pi i n z}$$

converging at all points of this strip, where

$$a_n = e^{\pi n(a+b)} \int_{x=0}^1 e^{-2\pi i n x} f\left(x + \frac{a+b}{2}i\right) dx.$$

(Suggestion: Consider the mapping  $w = e^{2\pi i z}$ . Show that  $f(z)$  determines an analytic function of  $w$  in the appropriate region, and use the two-sided expansion. Indicate where the hypothesis  $f(z + 1) = f(z)$  is needed.)

5. Find all singular points (either in the finite complex plane  $\mathbb{C}$  or at  $\infty$ ) of the following functions, and for each one indicate whether it is nonisolated, essential, a pole, or removable.

- (a)  $f(z) = 1/[(2z + 1)(z - 1)]$ ;
- (b)  $f(z) = 1/(e^z - 1)$ ;
- (c)  $f(z) = \pi/\sin(\pi/z)$ .