

Antiderivatives and Initial Value Problems

10/19/2011

Warm up

If $\frac{d}{dx}f(x) = 2x$, what is $f(x)$?

Can you think of another function that $f(x)$ could be?

If $\frac{d}{dx}f(x) = 3x^2 + 1$, what is $f(x)$?

Can you think of another function that $f(x)$ could be?

Definition

An **antiderivative** of a function f on an interval I is another function F such that $F'(x) = f(x)$ for all $x \in I$.

Definition

An **antiderivative** of a function f on an interval I is another function F such that $F'(x) = f(x)$ for all $x \in I$.

Examples:

1. An antiderivative of $f(x) = 2x$ is $F(x) = x^2$.

Definition

An **antiderivative** of a function f on an interval I is another function F such that $F'(x) = f(x)$ for all $x \in I$.

Examples:

1. An antiderivative of $f(x) = 2x$ is $F(x) = x^2$.
2. Another antiderivative of $f(x) = 2x$ is $F(x) = x^2 + 1$.

Definition

An **antiderivative** of a function f on an interval I is another function F such that $F'(x) = f(x)$ for all $x \in I$.

Examples:

1. An antiderivative of $f(x) = 2x$ is $F(x) = x^2$.
2. Another antiderivative of $f(x) = 2x$ is $F(x) = x^2 + 1$.
3. There are *lots* of antiderivatives of $f(x) = 2x$ which look like $F(x) = x^2 + c$.

Suppose that h is differentiable in an interval I ,
and $h'(x) = 0$ for all x in I .

Then h is a constant function!

i.e. $h(x) = C$ for all $x \in I$, where c is a constant.

So, if $F(x)$ is one antiderivative of $f(x)$, then any other antiderivative must be of the form $F(x) + c$.

Suppose that h is differentiable in an interval I , and $h'(x) = 0$ for all x in I .

Then h is a constant function!

i.e. $h(x) = C$ for all $x \in I$, where c is a constant.

So, if $F(x)$ is one antiderivative of $f(x)$, then any other antiderivative must be of the form $F(x) + c$.

Example: All of the antiderivatives of $f(x) = 2x$ look like

$$F(x) = x^2 + c$$

for some constant c .

Every function f that has at least one antiderivative F has **infinitely many** antiderivatives

$$F(x) + c.$$

Every function f that has at least one antiderivative F has **infinitely many** antiderivatives

$$F(x) + c.$$

We refer to $F(x) + c$ as the

general antiderivative or the **indefinite integral**

and denote it by

$$F(x) + c = \int f(x)dx.$$

Every function f that has at least one antiderivative F has **infinitely many** antiderivatives

$$F(x) + c.$$

We refer to $F(x) + c$ as the

general antiderivative or the **indefinite integral**

and denote it by

$$F(x) + c = \int f(x)dx.$$

Example:

$$\int 2x \, dx = x^2 + c.$$

Examples

$$\int x^2 dx = \frac{1}{3}x^3 + c, \quad \text{because} \quad \frac{d}{dx}\left(\frac{1}{3}x^3 + c\right) = \frac{1}{3} * 3x^2 = x^2$$

Examples

$$\int x^2 dx = \frac{1}{3}x^3 + c, \quad \text{because} \quad \frac{d}{dx}\left(\frac{1}{3}x^3 + c\right) = \frac{1}{3} * 3x^2 = x^2$$

$$\int x^3 dx = \frac{1}{4}x^4 + c, \quad \text{because} \quad \frac{d}{dx}\left(\frac{1}{4}x^4 + c\right) = \frac{1}{4} * 4x^3 = x^3$$

Examples

$$\int x^2 dx = \frac{1}{3}x^3 + c, \quad \text{because} \quad \frac{d}{dx}\left(\frac{1}{3}x^3 + c\right) = \frac{1}{3} * 3x^2 = x^2$$

$$\int x^3 dx = \frac{1}{4}x^4 + c, \quad \text{because} \quad \frac{d}{dx}\left(\frac{1}{4}x^4 + c\right) = \frac{1}{4} * 4x^3 = x^3$$

$$\int x^5 dx =$$

$$\int x^{-3} dx =$$

$$\int x^k dx =$$

$$\int x^k dx = \frac{1}{k+1} x^{k+1} + c \quad \text{if } k \neq -1$$

$$\int x^{-1} dx =$$

$$\int x^k dx = \frac{1}{k+1} x^{k+1} + c \quad \text{if } k \neq -1$$

$$\int x^{-1} dx = \ln(x) + c$$

$$\text{b/c } \frac{d}{dx} \ln(x) = x^{-1}$$

$$\int x^k dx = \frac{1}{k+1} x^{k+1} + c \quad \text{if } k \neq -1$$

$$\int x^{-1} dx = \ln(x) + c \quad \text{b/c } \frac{d}{dx} \ln(x) = x^{-1}$$

$$\int \sin(x) dx =$$

$$\int \cos(x) dx =$$

$$\int e^x dx =$$

$$\int \sec^2(x) dx =$$

Theorem (Opposite of sum and constant rules)

Suppose the functions f and g both have antiderivatives on the interval I . Then for any constants a and b , the function $af + bg$ has an antiderivative on I and

$$\int (a * f(x) + b * g(x)) dx = a \int f(x) dx + b \int g(x) dx$$

Remember differential equations?

We did

“Solve the differential equation $\frac{d}{dx}y = y.$ ”

Answer: $y = ce^x$

Remember differential equations?

We did

“Solve the differential equation $\frac{d}{dx}y = y$.”

Answer: $y = ce^x$

Finding an antiderivative can also be thought of as solving a differential equation:

“Solve the differential equation $\frac{d}{dx}y = x^2$.”

Answer: $y = \int x^2 dx = \frac{1}{3}x^3 + c$.

Remember differential equations?

We did

“Solve the differential equation $\frac{d}{dx}y = y$.”

Answer: $y = ce^x$

Finding an antiderivative can also be thought of as solving a differential equation:

“Solve the differential equation $\frac{d}{dx}y = x^2$.”

Answer: $y = \int x^2 dx = \frac{1}{3}x^3 + c$.

In general, equations that involve $x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots$ are called *differential equations*.

Remember differential equations?

We did

“Solve the differential equation $\frac{d}{dx}y = y$.”

Answer: $y = ce^x$

Finding an antiderivative can also be thought of as solving a differential equation:

“Solve the differential equation $\frac{d}{dx}y = x^2$.”

Answer: $y = \int x^2 dx = \frac{1}{3}x^3 + c$.

In general, equations that involve $x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots$ are called *differential equations*.

To solve: find a function $f(x)$ that satisfies the equation identically when substituted for the unknown function y , i.e. let $y = f(x)$.

Examples

Solve the differential equation $y' = 2x + \sin x$.

Solve the differential equation $\frac{d^2y}{dx^2} + y = 0$.

Example

Find a solution to the differential equation $\frac{d}{dx}y = x^2 + 1$ which *also* satisfies $y(2) = 8/3$.

Example


Find a solution to the differential equation $\frac{d}{dx}y = x^2 + 1$ which *also* satisfies $y(2) = 8/3$.

general solution: $y = \frac{1}{3}x^3 + x + c$
--

Example

Find a solution to the differential equation $\frac{d}{dx}y = x^2 + 1$ which *also* satisfies $y(2) = 8/3$.

$$\text{general solution: } y = \frac{1}{3}x^3 + x + c$$




Each color corresponds to a choice of c .

Example

Find a solution to the differential equation $\frac{d}{dx}y = x^2 + 1$ which *also* satisfies $y(2) = 8/3$.

$$\text{general solution: } y = \frac{1}{3}x^3 + x + c$$




Each color corresponds to a choice of c .

Example

Find a solution to the differential equation $\frac{d}{dx}y = x^2 + 1$ which *also* satisfies $y(2) = 8/3$.

$$\text{general solution: } y = \frac{1}{3}x^3 + x + c$$




Each color corresponds to a choice of c .

Example

Find a solution to the differential equation $\frac{d}{dx}y = x^2 + 1$ which *also* satisfies $y(2) = 8/3$.

$$\text{general solution: } y = \frac{1}{3}x^3 + x + c$$



Each color corresponds to a choice of c .


Red curve is the *particular* solution.

Example

Find a solution to the differential equation $\frac{d}{dx}y = x^2 + 1$ which *also* satisfies $y(2) = 8/3$.

general solution:	$y = \frac{1}{3}x^3 + x + c$
-------------------	------------------------------

particular solution:	$y = \frac{1}{3}x^3 + x - 2$
----------------------	------------------------------



Each color corresponds to a choice of c .

Red curve is the *particular* solution.

Definition

An **initial-value problem** is a differential equation together with enough additional conditions to specify the constants of integration that appear in the general solution.

The **particular solution of the problem** is then a function that satisfies both the differential equation and also the additional conditions.

Solve the initial value problem

$$\frac{dy}{dx} = 2x + \sin(x)$$

subject to $y(0) = 0$.

Solve the initial value problem

$$\frac{dy}{dx} = 2x + \sin(x)$$

subject to $y(0) = 0$.


general solution: $y = x^2 - \cos(x) + c$

Solve the initial value problem

$$\frac{dy}{dx} = 2x + \sin(x)$$

subject to $y(0) = 0$.

general solution: $y = x^2 - \cos(x) + c$




Solve the initial value problem

$$\frac{dy}{dx} = 2x + \sin(x)$$

subject to $y(0) = 0$.

general solution: $y = x^2 - \cos(x) + c$




Solve the initial value problem

$$\frac{dy}{dx} = 2x + \sin(x)$$

subject to $y(0) = 0$.

general solution: $y = x^2 - \cos(x) + c$




Solve the initial value problem

$$\frac{dy}{dx} = 2x + \sin(x)$$

subject to $y(0) = 0$.

general solution: $y = x^2 - \cos(x) + c$




Algebraically: get a particular solution by solving
 $0 = y(0) = (0)^2 - \cos(0) + c = -1 + c$ (for c)

Solve the initial value problem

$$\frac{dy}{dx} = 2x + \sin(x)$$

subject to $y(0) = 0$.

general solution: $y = x^2 - \cos(x) + c$



Algebraically: get a particular solution by solving
 $0 = y(0) = (0)^2 - \cos(0) + c = -1 + c$ (for c)


$c = 1,$ so $y = x^2 - \cos(x) + 1.$

Solve the initial value problem

$$\frac{dy}{dx} = 2x + \sin(x)$$

subject to $y(0) = 0$.

general solution: $y = x^2 - \cos(x) + c$



Algebraically: get a particular solution by solving
 $0 = y(0) = (0)^2 - \cos(0) + c = -1 + c$ (for c)

$c = 1,$ so $y = x^2 - \cos(x) + 1.$

Solve the initial-value problem $y'' = \cos x$, $y'(\frac{\pi}{2}) = 2$, $y(\frac{\pi}{2}) = 3\pi$.

Solve the initial-value problem $y'' = \cos x$, $y'(\frac{\pi}{2}) = 2$, $y(\frac{\pi}{2}) = 3\pi$.

Step 1: Calculate the antiderivative of $\cos(x)$ to find the general solution for y' .

Step 2: Plug in the values $y'(\frac{\pi}{2}) = 2$ to calculate c .

Step 3: Write down the *particular* solution for y' .

Step 4: Calculate the antiderivative of your particular solution in Step 3 to find the *general solution* for y .

Step 5: Plug in the values $y(\frac{\pi}{2}) = 3\pi$ to solve for the new constant.

Step 6: Write down the *particular* solution for y .

Word problem:

An object dropped from a cliff has acceleration $a = -9.8 \text{ m/sec}^2$ under the influence of gravity. What is the function $s(t)$ that models its height at time t ?

Initial value problem:

Solve

$$\frac{d^2s}{dt^2} = -9.8, \quad s(0) = s_0, \quad s'(0) = 0.$$

Word problem:

An object **dropped** from a cliff has acceleration $a = -9.8 \text{ m/sec}^2$ under the influence of gravity. What is the function $s(t)$ that models its height at time t ?

Initial value problem:

Solve

$$\frac{d^2s}{dt^2} = -9.8, \quad s(0) = s_0, \quad s'(0) = 0.$$

Word problem:

An object dropped from a cliff has acceleration $a = -9.8 \text{ m/sec}^2$ under the influence of gravity. What is the function $s(t)$ that models its height at time t ?

Initial value problem:

Solve

$$\frac{d^2s}{dt^2} = -9.8, \quad s(0) = s_0, \quad s'(0) = 0.$$

Word problem:

Suppose that a baseball is thrown upward from the roof of a 100 meter high building. It hits the street below eight seconds later. What was the initial velocity of the baseball, and how high did it rise above the street before beginning its descent?

Initial value problem:

Solve

$$\frac{d^2s}{dt^2} = -9.8, \quad s(0) = 100, \quad s(8) = 0.$$

Use your solution to

- (1) calculate $s'(0)$, and
- (2) solve $s'(t_1) = 0$ for t_1 and calculate $s(t_1)$.

Word problem:

Suppose that a baseball is thrown upward from the roof of a **100 meter high building**. It hits the street below eight seconds later. What was the initial velocity of the baseball, and how high did it rise above the street before beginning its descent?

Initial value problem:

Solve

$$\frac{d^2s}{dt^2} = -9.8, \quad s(0) = 100, \quad s(8) = 0.$$

Use your solution to

- (1) calculate $s'(0)$, and
- (2) solve $s'(t_1) = 0$ for t_1 and calculate $s(t_1)$.

Word problem:

Suppose that a baseball is thrown upward from the roof of a 100 meter high building. It **hits the street below eight seconds later**. What was the initial velocity of the baseball, and how high did it rise above the street before beginning its descent?

Initial value problem:

Solve

$$\frac{d^2s}{dt^2} = -9.8, \quad s(0) = 100, \quad s(8) = 0.$$

Use your solution to

- (1) calculate $s'(0)$, and
- (2) solve $s'(t_1) = 0$ for t_1 and calculate $s(t_1)$.

Word problem:

Suppose that a baseball is thrown upward from the roof of a 100 meter high building. It hits the street below eight seconds later. What was the **initial velocity** of the baseball, and how high did it rise above the street before beginning its descent?

Initial value problem:

Solve

$$\frac{d^2s}{dt^2} = -9.8, \quad s(0) = 100, \quad s(8) = 0.$$

Use your solution to

- (1) **calculate** $s'(0)$, and
- (2) solve $s'(t_1) = 0$ for t_1 and calculate $s(t_1)$.

Word problem:

Suppose that a baseball is thrown upward from the roof of a 100 meter high building. It hits the street below eight seconds later. What was the initial velocity of the baseball, and **how high did it rise** above the street before beginning its descent?

Initial value problem:

Solve

$$\frac{d^2s}{dt^2} = -9.8, \quad s(0) = 100, \quad s(8) = 0.$$

Use your solution to

- (1) calculate $s'(0)$, and
- (2) solve $s'(t_1) = 0$ for t_1 and **calculate $s(t_1)$.**