


# Curve Sketching

11/2/2011


## Warm up

Below are pictured six functions:  $f, f', f'', g, g',$  and  $g''$ . Pick out the two functions that could be  $f$  and  $g$ , and match them to their first and second derivatives, respectively.


(a)




(b)




(c)




(e)



(f)



(g)



## Review: Monotonicity of functions on intervals

Suppose that the function  $f$  is defined on an interval  $I$ , and let  $x_1$  and  $x_2$  denote points in  $I$ :

1.  $f$  is increasing on  $I$  if  $f(x_1) < f(x_2)$  whenever  $x_1 < x_2$ .
2.  $f$  is decreasing on  $I$  if  $f(x_1) > f(x_2)$  whenever  $x_1 < x_2$ .
3.  $f$  is constant on  $I$  if  $f(x_1) = f(x_2)$  for any  $x_1, x_2$  in  $I$ .

## Review: Testing monotonicity via derivatives


**Recall:** The derivative function  $f'(x)$  tells us the slope of the tangent line to the graph of the function  $f$  at the point  $(x, f(x))$ .

### Theorem (Increasing/Decreasing Test)

Suppose that  $f$  is continuous on  $[a, b]$  and differentiable on the open interval  $(a, b)$ . Then

1. If  $f'(x) > 0$  for every  $x$  in  $(a, b)$ , then  $f$  is increasing on  $[a, b]$ .
2. If  $f'(x) < 0$  for every  $x$  in  $(a, b)$ , then  $f$  is decreasing on  $[a, b]$ .
3. If  $f'(x) = 0$  for every  $x$  in  $(a, b)$ , then  $f$  is constant on  $[a, b]$ .

What it looks like:



## Theorem (Extreme Value Theorem)

If  $f$  is continuous on a closed interval  $[a, b]$ , then

1. there is a point  $c_1$  in the interval where  $f$  assumes its maximum value, i.e.  $f(x) \leq f(c_1)$  for all  $x$  in  $[a, b]$ , and
2. there is a point  $c_2$  in the interval where  $f$  assumes its maximal value, i.e.  $f(x) \geq f(c_2)$  for all  $x$  in  $[a, b]$ .


Finding minima and maxima is all about *optimizing* a function. So how do we find these values?

# Finding Extreme Values with Derivatives

## Theorem

If  $f$  is continuous in an open interval  $(a, b)$  and achieves a maximum (or minimum) value at a point  $c$  in  $(a, b)$  where  $f'(c)$  exists, then either  $f'(c)$  is not defined or  $f'(c) = 0$ .


**Big Idea:** if  $f'(c)$  exists, and is not equal to 0, then  $f(x)$  is either increasing or decreasing on both sides of  $c$ , so  $f(c)$  could not be a min or a max.



**Definition:** A point  $x = c$  where  $f'(c) = 0$  or where  $f'(c)$  doesn't exist is called a **critical point**. If  $f'(c)$  is undefined,  $c$  is also called a **singular point**.

**Warning:** Not all critical points are local minima or maxima:

**Example:** If  $f(x) = x^3$ , then  $f'(x) = 3x^2$ , and so  $f'(0) = 0$ :





## Strategy for closed bounded intervals

1. Calculate  $f'(x)$ .
2. Find where  $f'(x)$  is 0 or undefined on  $[a, b]$  (critical/singular points).
3. Evaluate  $f(x)$  at the critical and singular points, and at endpoints. The largest (reps. smallest) value among these is the maximum (reps. minimum).

## Strategy for closed bounded intervals

1. Calculate  $f'(x)$ .
2. Find where  $f'(x)$  is 0 or undefined on  $[a, b]$  (critical/singular points).
3. Evaluate  $f(x)$  at the critical and singular points, and at endpoints. The largest (reps. smallest) value among these is the maximum (reps. minimum).

**Example:** Let  $f(x) = x^3 - 3x$ . What are the min/max values on the interval  $[0, 2]$ .

## Strategy for closed bounded intervals

1. Calculate  $f'(x)$ .
2. Find where  $f'(x)$  is 0 or undefined on  $[a, b]$  (critical/singular points).
3. Evaluate  $f(x)$  at the critical and singular points, and at endpoints. The largest (reps. smallest) value among these is the maximum (reps. minimum).

**Example:** Let  $f(x) = x^3 - 3x$ . What are the min/max values on the interval  $[0, 2]$ .

$$f'(x) = 3x^2 - 3 = 3(x + 1)(x - 1)$$

## Strategy for closed bounded intervals

1. Calculate  $f'(x)$ .
2. Find where  $f'(x)$  is 0 or undefined on  $[a, b]$  (critical/singular points).
3. Evaluate  $f(x)$  at the critical and singular points, and at endpoints. The largest (reps. smallest) value among these is the maximum (reps. minimum).

**Example:** Let  $f(x) = x^3 - 3x$ . What are the min/max values on the interval  $[0, 2]$ .

$$f'(x) = 3x^2 - 3 = 3(x + 1)(x - 1)$$

So  $f'(x) = 0$  if  $x = -1$  or  $1$ .

## Strategy for closed bounded intervals

1. Calculate  $f'(x)$ .
2. Find where  $f'(x)$  is 0 or undefined on  $[a, b]$  (critical/singular points).
3. Evaluate  $f(x)$  at the critical and singular points, and at endpoints. The largest (reps. smallest) value among these is the maximum (reps. minimum).

**Example:** Let  $f(x) = x^3 - 3x$ . What are the min/max values on the interval  $[0, 2]$ .

$$f'(x) = 3x^2 - 3 = 3(x + 1)(x - 1)$$

So  $f'(x) = 0$  if  $x = -1$  or  $\boxed{1}$ .

## Strategy for closed bounded intervals

1. Calculate  $f'(x)$ .
2. Find where  $f'(x)$  is 0 or undefined on  $[a, b]$  (critical/singular points).
3. Evaluate  $f(x)$  at the critical and singular points, and at endpoints. The largest (reps. smallest) value among these is the maximum (reps. minimum).

**Example:** Let  $f(x) = x^3 - 3x$ . What are the min/max values on the interval  $[0, 2]$ .

$$f'(x) = 3x^2 - 3 = 3(x + 1)(x - 1)$$

So  $f'(x) = 0$  if  $x = -1$  or  $\boxed{1}$ .

$x$	$f(x)$	
1	-2	critical points
0	0	end points
2	2	

## Strategy for closed bounded intervals


1. Calculate  $f'(x)$ .
2. Find where  $f'(x)$  is 0 or undefined on  $[a, b]$  (critical/singular points).
3. Evaluate  $f(x)$  at the critical and singular points, and at endpoints. The largest (reps. smallest) value among these is the maximum (reps. minimum).

**Example:** Let  $f(x) = x^3 - 3x$ . What are the min/max values on the interval  $[0, 2]$ .

$$f'(x) = 3x^2 - 3 = 3(x + 1)(x - 1)$$

So  $f'(x) = 0$  if  $x = -1$  or  $\boxed{1}$ .

$x$	$f(x)$	
1	-2	critical points
0	0	end points
2	2	




# First Derivative Test


Finding local extrema can be useful for sketching curves.

Let  $c$  be a critical/singular point of a function  $y = f(x)$  that is continuous on an open interval  $I = (a, b)$  containing  $c$ . If  $f$  is differentiable on the interval (except possibly at the singular point  $x = c$ ) then the value  $f(c)$  can be classified as follows:

1. If  $f'(x)$  changes from positive to negative at  $x = c$ , then  $f(c)$  is a local maximum.



2. If  $f'(x)$  changes from negative to positive at  $x = c$ , then  $f(c)$  is a local minimum.



3. If  $f'(x)$  doesn't change sign, then it's neither a min or a max.



## Example

Find the local extrema of  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$  over the whole real line.

## Example

Find the local extrema of  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$  over the whole real line.

Calculate  $f'(x)$ :

## Example

Find the local extrema of  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$  over the whole real line.

Calculate  $f'(x)$ :


$$f'(x) = 12x^3 + 12x^2 - 2x - 2 = 12(x + 1)(x - 1/\sqrt{6})(x + 1/\sqrt{6})$$

## Example

Find the local extrema of  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$  over the whole real line.

Calculate  $f'(x)$ :

$$f'(x) = 12x^3 + 12x^2 - 2x - 2 = 12(x + 1)(x - 1/\sqrt{6})(x + 1/\sqrt{6})$$




## Example

Find the local extrema of  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$  over the whole real line.

Calculate  $f'(x)$ :

$$f'(x) = 12x^3 + 12x^2 - 2x - 2 = 12(x + 1)(x - 1/\sqrt{6})(x + 1/\sqrt{6})$$




## Example

Find the local extrema of  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$  over the whole real line.

Calculate  $f'(x)$ :

$$f'(x) = 12x^3 + 12x^2 - 2x - 2 = 12(x + 1)(x - 1/\sqrt{6})(x + 1/\sqrt{6})$$




## Example

Find the local extrema of  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$  over the whole real line.

Calculate  $f'(x)$ :

$$f'(x) = 12x^3 + 12x^2 - 2x - 2 = 12(x + 1)(x - 1/\sqrt{6})(x + 1/\sqrt{6})$$




## Example

Find the local extrema of  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$  over the whole real line.

Calculate  $f'(x)$ :

$$f'(x) = 12x^3 + 12x^2 - 2x - 2 = 12(x + 1)(x - 1/\sqrt{6})(x + 1/\sqrt{6})$$







## Example

Find the local extrema of  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$  over the whole real line.

Calculate  $f'(x)$ :

$$f'(x) = 12x^3 + 12x^2 - 2x - 2 = 12(x + 1)(x - 1/\sqrt{6})(x + 1/\sqrt{6})$$




## Example

Find the local extrema of  $f(x) = \frac{x^4 + 1}{x^2}$  over the whole real line.


[Hint: find a common denominator after taking a derivative.]

# Concavity

Q. How can we measure when a function is concave up or down?




Concave up  
 $f'(x)$  is increasing



Concave down  
 $f'(x)$  is decreasing


# Concavity

Q. How can we measure when a function is concave up or down?



Concave up  
 $f'(x)$  is increasing

$$f''(x) > 0$$






Concave down  
 $f'(x)$  is decreasing

$$f''(x) < 0$$

# Concavity and Inflection Points


**Definition:** The function  $f$  has an **inflection point** at the point  $x = c$  if  $f'(c)$  exists and the concavity changes at  $x = c$  from up to down or vice versa.



Back to the example  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$

Find the inflection points of  $f(x)$ , and where  $f(x)$  is concave up or down.


We calculated  $f'(x) = 12x^3 + 12x^2 - 2x - 2$ .



Back to the example  $f(x) = 3x^4 + 4x^3 - x^2 - 2x$

Find the inflection points of  $f(x)$ , and where  $f(x)$  is concave up or down.

We calculated  $f'(x) = 12x^3 + 12x^2 - 2x - 2$ .




So

$$f''(x) = 36x^2 + 24x - 2 = (6x - (\sqrt{6} - 2))(6x + \sqrt{6} + 2)$$

## Back to the example $f(x) = 3x^4 + 4x^3 - x^2 - 2x$


Find the inflection points of  $f(x)$ , and where  $f(x)$  is concave up or down.

We calculated  $f'(x) = 12x^3 + 12x^2 - 2x - 2$ .



So

$$f''(x) = 36x^2 + 24x - 2 = (6x - (\sqrt{6} - 2))(6x + \sqrt{6} + 2)$$






## Back to the example $f(x) = 3x^4 + 4x^3 - x^2 - 2x$


Find the inflection points of  $f(x)$ , and where  $f(x)$  is concave up or down.

We calculated  $f'(x) = 12x^3 + 12x^2 - 2x - 2$ .



So


$$f''(x) = 36x^2 + 24x - 2 = (6x - (\sqrt{6} - 2))(6x + \sqrt{6} + 2)$$



## Back to the example $f(x) = 3x^4 + 4x^3 - x^2 - 2x$


Find the inflection points of  $f(x)$ , and where  $f(x)$  is concave up or down.

We calculated  $f'(x) = 12x^3 + 12x^2 - 2x - 2$ .






So

$$f''(x) = 36x^2 + 24x - 2 = (6x - (\sqrt{6} - 2))(6x + \sqrt{6} + 2)$$



# Putting it together





# The second derivative test

## Theorem

Let  $f$  be a function whose second derivative exists on an interval  $I$  containing  $x_0$ .

1. If  $f'(x_0) = 0$  and  $f''(x_0) > 0$ , then  $f(x_0)$  is a local minimum.
2. If  $f'(x_0) = 0$  and  $f''(x_0) < 0$ , then  $f(x_0)$  is a local maximum.
3. If  $f'(x_0) = 0$  and  $f''(x_0) \leq 0$ , then the test **fails**, use the first derivative test to decide.



Sketch graphs of the following functions:

1.  $f(x) = -3x^5 + 5x^3$ .

2.  $f(x) = \frac{x^2 - 1}{x^2 + 1}$

### Instructions:

**Step 1** Find any places where  $f(x)$  is 0 or undefined.

**Step 2** Calculate  $f'(x)$  and find critical/singular points.

**Step 3** Classify where  $f'(x)$  is positive/negative, and therefore where  $f(x)$  is increasing/decreasing.

**Step 4** Calculate  $f''(x)$ , and find where it's 0 or undefined.

**Step 5** Classify where  $f''(x)$  is positive/negative, and therefore where  $f(x)$  is concave up/down.

**Step 6** Calculate  $\lim_{x \rightarrow \infty} f(x)$  and  $\lim_{x \rightarrow -\infty} f(x)$  to see what the tails are doing.

Hint for 2: Always simplify as a fraction of polynomials after taking a derivative.