

Using implicit differentiation for good: Inverse functions.

Warmup: Calculate $\frac{dy}{dx}$ if

1. $e^y = xy$

2. $\cos(y) = x + y$

Every time:

(1) Take $\frac{d}{dx}$ of both sides.

(2) Add and subtract to get the $\frac{dy}{dx}$ terms on one side and everything else on the other.

(3) Factor out $\frac{dy}{dx}$ and divide both sides by its coefficient.

Using implicit differentiation for good: Inverse functions.

Warmup: Calculate $\frac{dy}{dx}$ if

1. $e^y = xy$

Take $\frac{d}{dx}$ of both sides to find $e^y \frac{dy}{dx} = x \frac{dy}{dx} + y$.

So

$$y = e^y \frac{dy}{dx} - x \frac{dy}{dx} = \frac{dy}{dx} (e^y - x), \quad \text{implying} \quad \boxed{\frac{dy}{dx} = \frac{y}{e^y - x}}$$

2. $\cos(y) = x + y$

Take $\frac{d}{dx}$ as before: $-\sin(y) \frac{dy}{dx} = 1 + \frac{dy}{dx}$. So

$$\frac{dy}{dx} (\sin(y) + 1) = -1, \quad \text{and so} \quad \boxed{\frac{dy}{dx} = \frac{-1}{\sin(y) + 1}}$$

Every time:

(1) Take $\frac{d}{dx}$ of both sides.

(2) Add and subtract to get the $\frac{dy}{dx}$ terms on one side and everything else on the other.


(3) Factor out $\frac{dy}{dx}$ and divide both sides by its coefficient.

The Derivative of $y = \ln x$

Remember:

- (1) $y = e^x$ has a slope through the point $(0,1)$ of 1.
- (2) The natural log is the **inverse** function of e^x , so

$$y = \ln x \quad \Leftrightarrow \quad e^y = x$$



The Derivative of $y = \ln x$

To find the derivative of $\ln(x)$, use implicit differentiation!

The Derivative of $y = \ln x$

To find the derivative of $\ln(x)$, use implicit differentiation!

Rewrite

$$y = \ln x \quad \text{as} \quad e^y = x$$

The Derivative of $y = \ln x$

To find the derivative of $\ln(x)$, use implicit differentiation!

Rewrite

$$y = \ln x \quad \text{as} \quad e^y = x$$

Take a derivative of both sides of $e^y = x$ to get

$$\frac{dy}{dx} e^y = 1 \quad \text{so}$$

The Derivative of $y = \ln x$

To find the derivative of $\ln(x)$, use implicit differentiation!

Rewrite

$$y = \ln x \quad \text{as} \quad e^y = x$$

Take a derivative of both sides of $e^y = x$ to get

$$\frac{dy}{dx} e^y = 1 \quad \text{so} \quad \frac{dy}{dx} = \frac{1}{e^y}$$

The Derivative of $y = \ln x$

To find the derivative of $\ln(x)$, use implicit differentiation!

Rewrite

$$y = \ln x \quad \text{as} \quad e^y = x$$

Take a derivative of both sides of $e^y = x$ to get

$$\frac{dy}{dx} e^y = 1 \quad \text{so} \quad \frac{dy}{dx} = \frac{1}{e^y}$$

Problem: We asked “what is the derivative of $\ln(x)$?” and got back an answer with y in it!

The Derivative of $y = \ln x$

To find the derivative of $\ln(x)$, use implicit differentiation!

Rewrite

$$y = \ln x \quad \text{as} \quad e^y = x$$

Take a derivative of both sides of $e^y = x$ to get

$$\frac{dy}{dx} e^y = 1 \quad \text{so} \quad \frac{dy}{dx} = \frac{1}{e^y}$$

Problem: We asked “what is the derivative of $\ln(x)$?” and got back and answer with y in it!

Solution: Substitute back!

$$\frac{dy}{dx} = \frac{1}{e^y} = \frac{1}{e^{\ln(x)}}$$

The Derivative of $y = \ln x$

To find the derivative of $\ln(x)$, use implicit differentiation!

Rewrite

$$y = \ln x \quad \text{as} \quad e^y = x$$

Take a derivative of both sides of $e^y = x$ to get

$$\frac{dy}{dx} e^y = 1 \quad \text{so} \quad \frac{dy}{dx} = \frac{1}{e^y}$$

Problem: We asked “what is the derivative of $\ln(x)$?” and got back an answer with y in it!

Solution: Substitute back!

$$\frac{dy}{dx} = \frac{1}{e^y} = \frac{1}{e^{\ln(x)}} = \frac{1}{x}$$

The Derivative of $y = \ln x$

To find the derivative of $\ln(x)$, use implicit differentiation!

Rewrite

$$y = \ln x \quad \text{as} \quad e^y = x$$

Take a derivative of both sides of $e^y = x$ to get

$$\frac{dy}{dx} e^y = 1 \quad \text{so} \quad \frac{dy}{dx} = \frac{1}{e^y}$$

Problem: We asked “what is the derivative of $\ln(x)$?” and got back and answer with y in it!

Solution: Substitute back!


$$\frac{dy}{dx} = \frac{1}{e^y} = \frac{1}{e^{\ln(x)}} = \frac{1}{x}$$

$$\boxed{\frac{d}{dx} \ln(x) = \frac{1}{x}}$$


Does it make sense?

$$\frac{d}{dx} \ln(x) = \frac{1}{x}$$

$$f(x) = \ln(x)$$



$$f(x) = \frac{1}{x}$$



Examples

Calculate

1. $\frac{d}{dx} \ln x^2$

2. $\frac{d}{dx} \ln(\sin(x^2))$

3. $\frac{d}{dx} \log_3(x)$

[hint: $\log_a x = \frac{\ln x}{\ln a}$]

Examples

Calculate

$$1. \frac{d}{dx} \ln x^2 = \frac{2x}{x^2} = \frac{2}{x}$$

$$2. \frac{d}{dx} \ln(\sin(x^2)) = \frac{2x \cos(x^2)}{\sin(x^2)}$$

$$3. \frac{d}{dx} \log_3(x) = \frac{1}{x \ln(3)}$$

$$[\text{hint: } \log_a x = \frac{\ln x}{\ln a}]$$

Examples

Calculate

$$1. \frac{d}{dx} \ln x^2 = \frac{2x}{x^2} = \frac{2}{x}$$

$$2. \frac{d}{dx} \ln(\sin(x^2)) = \frac{2x \cos(x^2)}{\sin(x^2)}$$

$$3. \frac{d}{dx} \log_3(x) = \frac{1}{x \ln(3)}$$

$$[\text{hint: } \log_a x = \frac{\ln x}{\ln a}]$$

Notice, every time:

$$\frac{d}{dx} \ln(f(x)) = \frac{f'(x)}{f(x)}$$

Quick tip: Logarithmic differentiation

Example: Calculate $\frac{dy}{dx}$ if $y = x^{\sin(x)}$

Quick tip: Logarithmic differentiation

Example: Calculate $\frac{dy}{dx}$ if $y = x^{\sin(x)}$

Problem: Both the base and the exponent have the variable in them! So we can't use

$$\frac{d}{dx}x^a = ax^{a-1} \quad \text{or} \quad \frac{d}{dx}a^x = \ln(a)a^x.$$

Quick tip: Logarithmic differentiation

Example: Calculate $\frac{dy}{dx}$ if $y = x^{\sin(x)}$

Problem: Both the base and the exponent have the variable in them! So we can't use

$$\frac{d}{dx}x^a = ax^{a-1} \quad \text{or} \quad \frac{d}{dx}a^x = \ln(a)a^x.$$

Fix: Take the log of both sides and use implicit differentiation:

Quick tip: Logarithmic differentiation

Example: Calculate $\frac{dy}{dx}$ if $y = x^{\sin(x)}$

Problem: Both the base and the exponent have the variable in them! So we can't use

$$\frac{d}{dx}x^a = ax^{a-1} \quad \text{or} \quad \frac{d}{dx}a^x = \ln(a)a^x.$$

Fix: Take the log of both sides and use implicit differentiation:

$$\ln(y) = \ln(x^{\sin(x)})$$

Quick tip: Logarithmic differentiation

Example: Calculate $\frac{dy}{dx}$ if $y = x^{\sin(x)}$

Problem: Both the base and the exponent have the variable in them! So we can't use

$$\frac{d}{dx}x^a = ax^{a-1} \quad \text{or} \quad \frac{d}{dx}a^x = \ln(a)a^x.$$

Fix: Take the log of both sides and use implicit differentiation:

$$\ln(y) = \ln(x^{\sin(x)}) = \sin(x) * \ln(x) \quad (\text{using } \ln(a^b) = b \ln(a))$$

Quick tip: Logarithmic differentiation

Example: Calculate $\frac{dy}{dx}$ if $y = x^{\sin(x)}$

Problem: Both the base and the exponent have the variable in them! So we can't use

$$\frac{d}{dx}x^a = ax^{a-1} \quad \text{or} \quad \frac{d}{dx}a^x = \ln(a)a^x.$$

Fix: Take the log of both sides and use implicit differentiation:

$$\ln(y) = \ln(x^{\sin(x)}) = \sin(x) * \ln(x) \quad (\text{using } \ln(a^b) = b \ln(a))$$

Taking the derivative of both sides gives

$$\frac{1}{y} \frac{dy}{dx} = \cos(x) \ln(x) + \sin(x) \frac{1}{x}$$

Quick tip: Logarithmic differentiation

Example: Calculate $\frac{dy}{dx}$ if $y = x^{\sin(x)}$

Problem: Both the base and the exponent have the variable in them! So we can't use

$$\frac{d}{dx}x^a = ax^{a-1} \quad \text{or} \quad \frac{d}{dx}a^x = \ln(a)a^x.$$

Fix: Take the log of both sides and use implicit differentiation:

$$\ln(y) = \ln(x^{\sin(x)}) = \sin(x) * \ln(x) \quad (\text{using } \ln(a^b) = b \ln(a))$$

Taking the derivative of both sides gives

$$\frac{1}{y} \frac{dy}{dx} = \cos(x) \ln(x) + \sin(x) \frac{1}{x}$$

Then solving for $\frac{dy}{dx}$,

$$\frac{dy}{dx} = y \left(\cos(x) \ln(x) + \sin(x) \frac{1}{x} \right)$$

Quick tip: Logarithmic differentiation

Example: Calculate $\frac{dy}{dx}$ if $y = x^{\sin(x)}$

Problem: Both the base and the exponent have the variable in them! So we can't use

$$\frac{d}{dx}x^a = ax^{a-1} \quad \text{or} \quad \frac{d}{dx}a^x = \ln(a)a^x.$$

Fix: Take the log of both sides and use implicit differentiation:

$$\ln(y) = \ln(x^{\sin(x)}) = \sin(x) * \ln(x) \quad (\text{using } \ln(a^b) = b \ln(a))$$

Taking the derivative of both sides gives

$$\frac{1}{y} \frac{dy}{dx} = \cos(x) \ln(x) + \sin(x) \frac{1}{x}$$

Then solving for $\frac{dy}{dx}$,

$$\frac{dy}{dx} = y \left(\cos(x) \ln(x) + \sin(x) \frac{1}{x} \right) = \boxed{x^{\sin(x)} \left(\cos(x) \ln(x) + \sin(x) \frac{1}{x} \right)}.$$

Back to inverses

In the case where $y = \ln(x)$, we used the fact that $\ln(x) = f^{-1}(x)$, where $f(x) = e^x$, and got

$$\frac{d}{dx} \ln(x) = \frac{1}{e^{\ln(x)}}.$$

In general, calculating $\frac{d}{dx} f^{-1}(x)$:

Back to inverses

In the case where $y = \ln(x)$, we used the fact that $\ln(x) = f^{-1}(x)$, where $f(x) = e^x$, and got

$$\frac{d}{dx} \ln(x) = \frac{1}{e^{\ln(x)}}.$$

In general, calculating $\frac{d}{dx} f^{-1}(x)$:

(1) Rewrite $y = f^{-1}(x)$ as $f(y) = x$.

Back to inverses

In the case where $y = \ln(x)$, we used the fact that $\ln(x) = f^{-1}(x)$, where $f(x) = e^x$, and got

$$\frac{d}{dx} \ln(x) = \frac{1}{e^{\ln(x)}}.$$

In general, calculating $\frac{d}{dx} f^{-1}(x)$:

(1) Rewrite $y = f^{-1}(x)$ as $f(y) = x$.

(2) Use implicit differentiation:

$$f'(y) * \frac{dy}{dx} = 1 \quad \text{so} \quad \boxed{\frac{dy}{dx} = \frac{1}{f'(y)} = \frac{1}{f'(f^{-1}(x))}}.$$

Examples

Just to check, use the rule

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

to calculate

1. $\frac{d}{dx} \ln(x)$ (the inverse of e^x)

2. $\frac{d}{dx} \sqrt{x}$ (the inverse of x^2)

Examples

Just to check, use the rule

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

to calculate

1. $\frac{d}{dx} \ln(x)$ (the inverse of e^x)

In the notation above, $f^{-1}(x) = \ln(x)$ and $f(x) = e^x$.

2. $\frac{d}{dx} \sqrt{x}$ (the inverse of x^2)

Examples

Just to check, use the rule

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

to calculate

1. $\frac{d}{dx} \ln(x)$ (the inverse of e^x)

In the notation above, $f^{-1}(x) = \ln(x)$ and $f(x) = e^x$.

We'll also need $f'(x) = e^x$.

2. $\frac{d}{dx} \sqrt{x}$ (the inverse of x^2)

Examples

Just to check, use the rule

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

to calculate

1. $\frac{d}{dx} \ln(x)$ (the inverse of e^x)

In the notation above, $f^{-1}(x) = \ln(x)$ and $f(x) = e^x$.

We'll also need $f'(x) = e^x$. So

$$\frac{d}{dx} \ln(x) = \frac{1}{e^{\ln(x)}} \quad \text{☺}$$

2. $\frac{d}{dx} \sqrt{x}$ (the inverse of x^2)

Examples

Just to check, use the rule

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

to calculate

1. $\frac{d}{dx} \ln(x)$ (the inverse of e^x)

In the notation above, $f^{-1}(x) = \ln(x)$ and $f(x) = e^x$.

We'll also need $f'(x) = e^x$. So

$$\frac{d}{dx} \ln(x) = \frac{1}{e^{\ln(x)}} \quad \text{☺}$$

2. $\frac{d}{dx} \sqrt{x}$ (the inverse of x^2)

In the notation above, $f^{-1}(x) = \sqrt{x}$ and $f(x) = x^2$.

Examples

Just to check, use the rule

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

to calculate

1. $\frac{d}{dx} \ln(x)$ (the inverse of e^x)

In the notation above, $f^{-1}(x) = \ln(x)$ and $f(x) = e^x$.

We'll also need $f'(x) = e^x$. So

$$\frac{d}{dx} \ln(x) = \frac{1}{e^{\ln(x)}} \quad \text{☺}$$

2. $\frac{d}{dx} \sqrt{x}$ (the inverse of x^2)

In the notation above, $f^{-1}(x) = \sqrt{x}$ and $f(x) = x^2$.

We'll also need $f'(x) = 2x$.

Examples

Just to check, use the rule

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

to calculate

1. $\frac{d}{dx} \ln(x)$ (the inverse of e^x)

In the notation above, $f^{-1}(x) = \ln(x)$ and $f(x) = e^x$.

We'll also need $f'(x) = e^x$. So

$$\frac{d}{dx} \ln(x) = \frac{1}{e^{\ln(x)}} \quad \text{☺}$$

2. $\frac{d}{dx} \sqrt{x}$ (the inverse of x^2)

In the notation above, $f^{-1}(x) = \sqrt{x}$ and $f(x) = x^2$.

We'll also need $f'(x) = 2x$. So

$$\frac{d}{dx} \sqrt{x} = \frac{1}{2 * (\sqrt{x})} \quad \text{☺}$$


Recall inverse trig functions

Two notations:

$f(x)$	$f^{-1}(x)$
$\sin(x)$	$\sin^{-1}(x) = \arcsin(x)$
$\cos(x)$	$\cos^{-1}(x) = \arccos(x)$
$\tan(x)$	$\tan^{-1}(x) = \arctan(x)$
$\sec(x)$	$\sec^{-1}(x) = \operatorname{arcsec}(x)$
$\csc(x)$	$\csc^{-1}(x) = \operatorname{arccsc}(x)$
$\cot(x)$	$\cot^{-1}(x) = \operatorname{arccot}(x)$

In general:

$\text{arc}___(-)$ takes in a ratio and spits out an angle:




$$\cos(\theta) = a/c \quad \text{so} \quad \arccos(a/c) = \theta$$

$$\sin(\theta) = b/c \quad \text{so} \quad \arcsin(b/c) = \theta$$

$$\tan(\theta) = b/a \quad \text{so} \quad \arctan(b/a) = \theta$$

In general:

$\arcsin(\quad)$ takes in a ratio and spits out an angle:



$$\cos(\theta) = a/c \quad \text{so} \quad \arccos(a/c) = \theta$$

$$\sin(\theta) = b/c \quad \text{so} \quad \arcsin(b/c) = \theta$$

$$\tan(\theta) = b/a \quad \text{so} \quad \arctan(b/a) = \theta$$


Domain problems:

$$\sin(0) = 0, \quad \sin(\pi) = 0, \quad \sin(2\pi) = 0, \quad \sin(3\pi) = 0, \dots$$

So which is the right answer to $\arcsin(0)$, really?


Domain/range

$$y = \sin(x)$$



Domain/range


$$y = \sin(x)$$



Domain/range

$$y = \sin(x)$$


$$y = \arcsin(x)$$



Domain: $-1 \leq x \leq 1$

Domain/range


$$y = \arcsin(x)$$



$$\text{Domain: } -1 \leq x \leq 1$$

Domain/range

$$y = \arcsin(x)$$




Domain: $-1 \leq x \leq 1$

Range: $-\pi/2 \leq y \leq \pi/2$


Domain/range

$$y = \cos(x)$$



Domain/range


$$y = \cos(x)$$



Domain/range

$$y = \cos(x)$$


$$y = \arccos(x)$$



$$\text{Domain: } -1 \leq x \leq 1$$

Domain/range


$$y = \arccos(x)$$



$$\text{Domain: } -1 \leq x \leq 1$$

Domain/range

$$y = \arccos(x)$$




Domain: $-1 \leq x \leq 1$

Range: $0 \leq y \leq \pi$


Domain/range

$$y = \tan(x)$$



Domain/range


$$y = \tan(x)$$



Domain/range

$$y = \tan(x)$$


$$y = \arctan(x)$$



Domain: $-\infty \leq x \leq \infty$

Domain/range


$$y = \arctan(x)$$



Domain: $-\infty \leq x \leq \infty$

Domain/range

$$y = \arctan(x)$$




Domain: $-\infty \leq x \leq \infty$

Range: $-\pi/2 < y < \pi/2$


Domain/range

$$y = \sec(x)$$



Domain/range


$$y = \sec(x)$$



Domain/range

$$y = \sec(x)$$


$$y = \operatorname{arcsec}(x)$$



Domain: $x \leq -1$ and $1 \leq x$

Domain/range


$$y = \operatorname{arcsec}(x)$$



Domain: $x \leq -1$ and $1 \leq x$

Domain/range

$$y = \operatorname{arcsec}(x)$$




Domain: $x \leq -1$ and $1 \leq x$

Range: $0 \leq y \leq \pi$


Domain/range

$$y = \csc(x)$$



Domain/range


$$y = \csc(x)$$



Domain/range

$$y = \csc(x)$$


$$y = \operatorname{arccsc}(x)$$



Domain: $x \leq -1$ and $1 \leq x$

Domain/range


$$y = \operatorname{arccsc}(x)$$



Domain: $x \leq -1$ and $1 \leq x$

Domain/range

$$y = \operatorname{arccsc}(x)$$




Domain: $x \leq -1$ and $1 \leq x$

Range: $-\pi/2 \leq y \leq \pi/2$


Domain/range

$$y = \cot(x)$$



Domain/range


$$y = \cot(x)$$



Domain/range

$$y = \cot(x)$$


$$y = \operatorname{arccot}(x)$$



Domain: $-\infty \leq x \leq \infty$

Domain/range


$$y = \operatorname{arccot}(x)$$



$$\text{Domain: } -\infty \leq x \leq \infty$$

Domain/range

$$y = \operatorname{arccot}(x)$$




Domain: $-\infty \leq x \leq \infty$


Range: $0 < y < \pi$

Graphs


$\arcsin(x)$




$\arccos(x)$




$\arctan(x)$




$\operatorname{arcsec}(x)$



$\operatorname{arccsc}(x)$



$\operatorname{arccot}(x)$



Recall:

$f(x)$	$f'(x)$
$\sin(x)$	$\cos(x)$
$\cos(x)$	$-\sin(x)$
$\tan(x)$	$\sec^2(x)$
$\sec(x)$	$\sec(x)\tan(x)$
$\csc(x)$	$-\csc(x)\cot(x)$
$\cot(x)$	$-\csc^2(x)$

Back to Derivatives

Use implicit differentiation to calculate the derivatives of

1. $\arcsin(x)$
2. $\arctan(x)$

Use the rule

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

to check your answers, and then to calculate the derivatives of the other inverse trig functions:

1. $\frac{d}{dx} \arccos(x)$
2. $\frac{d}{dx} \operatorname{arcsec}(x)$
3. $\frac{d}{dx} \operatorname{arccsc}(x)$
4. $\frac{d}{dx} \operatorname{arccot}(x)$

Back to Derivatives

Use implicit differentiation to calculate the derivatives of

$$1. \arcsin(x) = \frac{1}{\cos(\arcsin(x))}$$

$$2. \arctan(x) = \frac{1}{\sec^2(\arctan(x))}$$

Use the rule

$$\frac{d}{dx} f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

to check your answers, and then to calculate the derivatives of the other inverse trig functions:

$$1. \frac{d}{dx} \arccos(x) = -\frac{1}{\sin(\arccos(x))}$$

$$2. \frac{d}{dx} \operatorname{arcsec}(x) = \frac{1}{\sec(\operatorname{arcsec}(x)) \tan(\operatorname{arcsec}(x))}$$

$$3. \frac{d}{dx} \operatorname{arccsc}(x) = -\frac{1}{\csc(\operatorname{arccsc}(x)) \cot(\operatorname{arccsc}(x))}$$

$$4. \frac{d}{dx} \operatorname{arccot}(x) = -\frac{1}{\csc^2(\operatorname{arccot}(x))}$$

Using implicit differentiation to calculate $\frac{d}{dx} \arcsin(x)$

$$\boxed{\text{If } y = \arcsin(x) \text{ then } x = \sin(y).}$$

Take $\frac{d}{dx}$ of both sides of $x = \sin(y)$:

$$\text{Left hand side: } \frac{d}{dx} x = 1$$


$$\text{Right hand side: } \frac{d}{dx} \sin(y) = \cos(y) * \frac{dy}{dx} = \cos(\arcsin(x)) * \frac{dy}{dx}$$

So

$$\boxed{\frac{dy}{dx} = \frac{1}{\cos(\arcsin(x))}.}$$

Simplifying $\cos(\arcsin(x))$


Call $\arcsin(x) = \theta$.



Simplifying $\cos(\arcsin(x))$

Call $\arcsin(x) = \theta$.


$$\sin(\theta) = x$$



Simplifying $\cos(\arcsin(x))$

Call $\arcsin(x) = \theta$.

$$\sin(\theta) = x$$




Key: This is a simple triangle to write down whose angle θ has $\sin(\theta) = x$

Simplifying $\cos(\arcsin(x))$

Call $\arcsin(x) = \theta$.


$$\sin(\theta) = x$$



Simplifying $\cos(\arcsin(x))$

Call $\arcsin(x) = \theta$.

$$\sin(\theta) = x$$




So $\cos(\theta) = \sqrt{1-x^2}/1$

Simplifying $\cos(\arcsin(x))$

Call $\arcsin(x) = \theta$.

$$\sin(\theta) = x$$




So $\cos(\arcsin(x)) = \sqrt{1-x^2}$

Simplifying $\cos(\arcsin(x))$

Call $\arcsin(x) = \theta$.

$$\sin(\theta) = x$$



$$\text{So } \cos(\arcsin(x)) = \sqrt{1-x^2}$$


$$\text{So } \frac{d}{dx} \arcsin(x) = \frac{1}{\cos(\arcsin(x))} = \frac{1}{\sqrt{1-x^2}}.$$

Calculating $\frac{d}{dx} \arctan(x)$.

We found that

$$\frac{d}{dx} \arctan(x) = \frac{1}{\sec^2(\arctan(x))} = \left(\frac{1}{\sec(\arctan(x))} \right)^2$$

Simplify this expression using




Calculating $\frac{d}{dx} \arctan(x)$.

We found that

$$\frac{d}{dx} \arctan(x) = \frac{1}{\sec^2(\arctan(x))} = \left(\frac{1}{\sec(\arctan(x))} \right)^2$$

Simplify this expression using



$$\frac{dy}{dx} = \left(\frac{1}{\sec(\arctan(x))} \right)^2 = \frac{1}{1+x^2}$$

To simplify the rest, use the triangles

